

CONSTRUCTION OF KÄHLER SURFACES WITH CONSTANT SCALAR CURVATURE

YANN ROLLIN AND MICHAEL SINGER

For Professor Paul Gauduchon on his 60th birthday

ABSTRACT. We present new constructions of Kähler metrics with constant scalar curvature on complex surfaces, in particular on certain del Pezzo surfaces. Some higher dimensional examples are provided as well.

1. INTRODUCTION

The aim of this note is to present a new construction of Kähler metrics of constant scalar curvature (CSC) on complex surfaces. In order to introduce our results, let us introduce the terms “positive CSC” to mean “constant positive scalar curvature”, “zero CSC” for “(constant) zero scalar curvature” and “negative CSC” for “constant negative scalar curvature”.

Our construction gives rise to many families of examples, but in this introduction we shall focus on

$$X_k := k\text{-fold blow-up of } \mathbb{CP}^1 \times \mathbb{CP}^1.$$

We note that if $k \geq 1$ then X_k can also be viewed as a $k + 1$ -fold blow-up of $\mathbb{CP}^2$. Of course, the above description of X_k does not fix its complex structure: this will depend on the location of the centres of the blow-ups. Our first result gives positive CSC Kähler metrics in a family of Kähler classes on X_k , for $k = 6, 7, 8$, and for certain choices of complex structure. We note that if $k \leq 7$ then X_k is Fano and the work of Tian [T] and others gives positive Kähler–Einstein metrics on X_k . Our result is new in that it produces CSC metrics on X_8 as well as CSC metrics on X_6 and X_7 in Kähler classes that are “arbitrarily far” from $c_1(X)$:

Date: December 2004.

First author partly supported by NSF grant # DMS-0305130.

Second author partly supported by a Leverhulme research fellowship and an EPSRC small grant.

Theorem A. *For $k = 6, 7, 8$, there exists a k -point blow-up X of $\mathbb{CP}^1 \times \mathbb{CP}^1$ with no non-trivial holomorphic vector field and the following properties. Let $F = \{x\} \times \mathbb{CP}^1$ be a generic rational curve of $\mathbb{CP}^1 \times \mathbb{CP}^1$. For every constant $c > 0$ and $\varepsilon > 0$, there exists a Kähler metric ω of strictly positive constant scalar curvature on X such that*

$$\left| \frac{[\omega] \cdot F}{\sqrt{[\omega]^2}} - c \right| \leq \varepsilon. \quad (1.1)$$

This theorem can be thought of as an exploration of the Kähler cone on certain del Pezzo surfaces. We are asking the question: which Kähler classes can be realized by Kähler metrics of strictly positive constant scalar curvature?

1.0.1. Remark. — For the KE metric on X_k , $k = 6, 7$, we have the number c takes the value

$$c^{\text{ein}} := \frac{[\omega^{\text{ein}}] \cdot F}{\sqrt{[\omega^{\text{ein}}]^2}} = \frac{c_1(X) \cdot F}{\sqrt{c_1^2(X)}} = \frac{2}{\sqrt{8-k}}.$$

1.0.2. Remark. — It is an observation of LeBrun (see, for example, [LS]) that the hyperplane

$$H = \{\alpha \in H^2(X, \mathbb{R}) : c_1 \cdot \alpha = 0\}$$

does not meet the Kähler cone of X_k unless $k \geq 9$. It follows that if $k \leq 8$, every Kähler class $[\omega]$ satisfies $c_1(X) \cdot [\omega] > 0$. Since this number represents, up to a factor of 4π , the integral of the scalar curvature over X , it follows that any CSC Kähler metric on X_k must be positive, whatever the Kähler class.

Now we turn to the case $k \geq 9$. In [RS], it was proved that there exist zero CSC Kähler metrics on X_k in this case (the same result was previously known for $k \geq 14$ [KLP]). In the next result, we obtain CSC Kähler metrics of either sign:

Theorem B. *There exists a 9-point blow-up X of $\mathbb{CP}^1 \times \mathbb{CP}^1$ with no non trivial holomorphic vector field and the following properties:*

- (i) *there is a scalar-flat Kähler metric on X ;*
- (ii) *for every $c \in \mathbb{R}$ and $\varepsilon > 0$, there exists a metric of constant scalar curvature ω on X such that*

$$\left| \frac{[\omega] \cdot c_1(X)}{\sqrt{[\omega]^2}} - c \right| \leq \varepsilon. \quad (1.2)$$

In addition, any further blow-up of X admits metrics with the same properties.

Part (i) of this Theorem was proved in [RS]. It plays in this discussion an analogous role to the one played by Kähler-Einstein metrics in Theorem A, and corresponds to $c = c^{\text{sfc}} = 0$. Again, the new content of this Theorem is that there are metrics of constant scalar curvature that represent Kähler classes arbitrarily far from the hyperplane H inside $H^2(X, \mathbb{R})$.

1.0.3. Remark. — Kähler metrics of constant scalar curvature admit a good perturbation theory: if the class $[\omega]$ is represented by a CSC Kähler metric, then the same is true for all sufficiently close Kähler classes, at least if X carries no non-zero holomorphic vector fields [LSc]. This applies to the KE metrics on X_6 and X_7 to give non-Einstein CSC metrics in all classes sufficiently close to $c_1(X)$. It also applies to give CSC Kähler metrics of either sign on X_9 , by perturbing a known zero CSC metric. It seems very difficult, however, to extend this perturbation theory to give more global results like Theorems A or B.

1.1. Strategy. The Theorems A and B will be deduced from an extension of the constructions invented in [RS] and the recent gluing theorem of Arezzo-Pacard [AP] for CSC Kähler metrics. The zero-CSC Kähler metrics in [RS] were obtained by resolving the singularities of a zero-CSC orbifold. The gluing theorem in that work applied only to produce zero-CSC Kähler metrics. The results of [AP] allow us to work with CSC orbifolds of non-zero scalar curvature, which are, however, constructed in almost exactly the same way as in [RS]. Indeed, as will be explained below, these orbifolds are twisted products of Riemann surfaces, where now the scalar curvature of the factors can be chosen arbitrarily.

1.1.1. Remark. — In §9(i) of their paper, Arezzo and Pacard make the following construction: take a copy of X_k with no holomorphic vector field and Tian's Kähler-Einstein metric (so $k \leq 7$). Their gluing theorem gives a positive CSC Kähler metric on the blow-up of X_k , that is to say, on X_{k+1} . If $k = 7$, then this proves Theorem A in the particular case $c = 2$, $k = 8$.

We should point out that in contrast to this approach, our construction does not use any result about the existence of Kähler-Einstein metrics on del Pezzo surfaces and may be considered to be more elementary for this reason.

1.1.2. Remark. — The gluing theorem of [AP] is not specific to complex dimension 2. Accordingly, in §6 we give some new but rather special higher-dimensional examples of negative CSC Kähler metrics.

Acknowledgments. We thank Claudio Arezzo, Denis Auroux and Frank Pacard for stimulating discussions.

2. THE FACTORY

We review the construction of scalar-flat Kähler orbifold metrics of [RS] and adapt it to the case of Kähler metrics of constant scalar curvature.

2.1. Orbifold Riemann surfaces. We start with a closed Riemann surface $\overline{\Sigma}$ of genus g with a finite set of orbifold points $P_1, P_2, \dots, P_k$, with local ramified cover of $q_1, q_2, \dots, q_k > 1$. Recall first the description of the fundamental group of the punctured Riemann surface $\Sigma = \overline{\Sigma} \setminus \{P_j\}$:

$$\pi_1(\Sigma) = \langle a_1, b_1, \dots, a_g, b_g, l_1, \dots, l_k : [a_1, b_1][a_2, b_2] \dots [a_g, b_g] l_1 \dots l_k = 1 \rangle$$

Here the a_j and b_j are standard generators of $\pi_1(\overline{\Sigma})$ and l_j is (the homotopy class of) a small loop around P_j . The orbifold fundamental group is defined by

$$\pi_1^{\text{orb}}(\overline{\Sigma}) = \pi_1(\Sigma)/G$$

where G is the normal subgroup of $\pi_1(\Sigma)$ generated by $l_1^{q_1}, l_2^{q_2}, \dots, l_k^{q_k}$.

The orbifold Euler characteristic is defined by

$$\chi^{\text{orb}}(\overline{\Sigma}) := \chi^{\text{top}}(\overline{\Sigma}) - \sum_{j=1}^k \left(1 - \frac{1}{q_j}\right).$$

The question of whether $\overline{\Sigma}$ carries a CSC orbifold Kähler metric was considered by Troyanov [Tr]. Let us call an orbifold Riemann surface “good” if its orbifold universal cover admits a compatible Kähler metric of CSC κ_1 , say. By the Gauss–Bonnet theorem (applied to the surface with boundary obtained by removing a small disc around each of the P_j), if $\overline{\Sigma}$ is good, then the sign of κ_1 is the same as the sign of $\chi^{\text{orb}}(\overline{\Sigma})$. However, not every orbifold Riemann surface is “good”: the *tear-drop*, which is S^2 with one orbifold point of any order ≥ 2 , is simply connected but admits no compatible metric of constant curvature.

The following summarizes the facts we shall need if the orbifold euler characteristic is non-positive:

Proposition 2.1.1. *The orbifold Riemann surface $\overline{\Sigma}$ is always good if*

$$\chi^{\text{orb}}(\overline{\Sigma}) \leq 0.$$

Such $\overline{\Sigma}$ carries no non-trivial holomorphic vector fields if $\chi^{\text{orb}}(\overline{\Sigma}) < 0$ or if $\chi^{\text{orb}}(\overline{\Sigma}) = 0$ and there is at least one orbifold point.

Proof. The existence of metrics of constant scalar curvature is contained in [Tr]. The statement about holomorphic vector fields follows either by lifting such a field to the universal cover or from the orbifold version of the Hopf index theorem: a holomorphic vector field must vanish at each of the orbifold points. Q.E.D.

For good orbifolds with positive orbifold euler characteristic, the universal cover must be $\mathbb{CP}^1$ and the orbifold fundamental group must be one of the finite subgroups of $SO(3)$:

Proposition 2.1.2. *Let $\overline{\Sigma}$ be a good orbifold with $\chi^{\text{orb}}(\overline{\Sigma}) > 0$. Then $\overline{\Sigma}$ is biholomorphic to $\mathbb{CP}^1/\Gamma$, where Γ is a finite subgroup of $SO(3)$. There are either 2 marked points of the same order $q \geq 2$ or 3 marked points with orders $\{2, 2, q\}$, $\{2, 3, 3\}$, $\{2, 3, 4\}$, $\{2, 3, 5\}$. The corresponding groups are cyclic of order q , dihedral, tetrahedral, octahedral, or icosahedral respectively.*

Furthermore, $\overline{\Sigma}$ carries no non-trivial holomorphic vector fields if and only if G is not cyclic.

Proof. We need only discuss the last part. The Euler vector field ($z\partial_z$ in affine coordinates) descends to the quotient if G is the cyclic group acting with fixed-points at $z = 0$ and $z = \infty$. On the other hand, no non-trivial holomorphic vector field on the sphere can have 3 zeros, so none of the other orbifolds can admit such vector fields. Q.E.D.

2.2. Desingularization of orbifold ruled surfaces. From now we on assume that $\overline{\Sigma}$ is a good orbifold Riemann surface and we endow $\overline{\Sigma}$ with an orbifold Kähler metric $\bar{g}^{\Sigma}$ of constant curvature κ_1 . Note that, just as for ordinary Riemann surfaces, we have

$$\overline{\Sigma} = \mathcal{U}/\pi_1^{\text{orb}}(\overline{\Sigma})$$

where the fundamental group acts by isometries on the universal cover $\mathcal{U}$ which is equal to $\mathbb{H}$, $\mathbb{E}$ or $\mathbb{CP}^1$, according as κ_1 is negative, zero, or positive.

Let g^{FS} be the Fubini-Study metric with curvature $\kappa_2 > 0$, on $\mathbb{CP}^1$. Let $\rho : \pi_1^{\text{orb}}(\overline{\Sigma}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$ be a homomorphism. Then $\pi_1^{\text{orb}}(\overline{\Sigma})$ acts isometrically on $\mathcal{U} \times \mathbb{CP}^1$. We deduce a Kähler metric $\bar{g}_\rho$ of constant scalar curvature $s = 2(\kappa_1 + \kappa_2)$ on the orbifold quotient

$$\overline{M}_\rho = \overline{\Sigma} \times_\rho \mathbb{CP}^1 = (\mathcal{U} \times \mathbb{CP}^1)/\pi_1^{\text{orb}}(\overline{\Sigma}).$$

The idea now is to apply the results of [AP] to obtain CSC Kähler metrics on the minimal resolution of singularities, $\widehat{M}_\rho$, say, of $\overline{M}_\rho$. We note that the constant s can be chosen arbitrarily if $\overline{\Sigma}$ is hyperbolic, but must be positive otherwise. Here is the full statement, which is parallel to Theorem D of [RS].

Theorem C. *Let $\overline{\Sigma}$ be a good compact orbifold Riemann surface as above, carrying no non-trivial holomorphic vector fields. Suppose that $\rho : \pi_1^{\text{orb}}(\overline{\Sigma}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$ is a homomorphism that is irreducible in the sense that the induced action of $\pi_1^{\text{orb}}(\overline{\Sigma})$ fixes no point of $\mathbb{CP}^1$. Equip $\overline{M}_\rho$ with a twisted product metric of CSC s as above.*

Then the minimal resolution $\widehat{M}_\rho \rightarrow \overline{M}_\rho$ carries a CSC Kähler metric with scalar curvature very close to s .

In addition, any further blow-up of $\widehat{M}_\rho$ carries a CSC Kähler metric with scalar curvature very close to s .

Proof. We have already equipped $\overline{M}_\rho$ with a Kähler orbifold metric of constant scalar curvature s . On the other hand, for each finite cyclic subgroup G of U_2 with the property that G acts freely on S^3 , there is an asymptotically locally euclidean zero-CSC Kähler metric on the minimal resolution [CS, RS]. If $s = 0$, we are in the situation of [RS], so we may as well assume that $s \neq 0$. We must check that the hypotheses of [AP] are satisfied, i.e. that $\overline{M}_\rho$ has no non-trivial holomorphic vector fields. But this follows as in [RS], for we have placed ourselves in the situation where the base $\overline{\Sigma}$ has no non-zero holomorphic vector fields.

Hence, we can indeed apply [AP] to obtain CSC Kähler metrics g on $\widehat{M}_\rho$. For sufficiently small choices of the gluing parameter, g is very close to g_ρ , and the scalar curvature of g will be very close to the number s .

To get the last statement, we again apply [AP], this time gluing a copy of the Burns metric at a smooth point. Q.E.D.

2.2.1. Remark. — The term “minimal” applied here is really unfortunate. According to its usual meaning in complex surface theory, it means that there are no rational curves of self-intersection -1 . In this case, however, the proper transform of any singular fibre of $\overline{M}_\rho$ is always a (-1) -curve! The term “minimal” is justified by the fact that we are taking the minimal resolution of each of the orbifold singularities separately. Only after doing so is it possible to blow down the (-1) -curves.

3. STABILITY OF PARABOLIC RULED SURFACES

In order to make the results of the previous section more useful, we need a way to generate representations $\rho : \pi_1^{\text{orb}}(\overline{\Sigma}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$. A way to do this was introduced in [RS], through the notion of a (stable) parabolic ruled surfaces. We recall the main definitions here:

A *geometrically ruled surface* $\widetilde{M}$ is by definition a minimal complex surface obtained as $\widetilde{M} = \mathbb{P}(E)$, where $E \rightarrow \widehat{\Sigma}$ is a holomorphic vector bundle

of rank 2 over a *smooth* Riemann surface $\widehat{\Sigma}$. The induced map $\pi : \widetilde{M} \rightarrow \widehat{\Sigma}$ is called the *ruling*.

A parabolic structure on $\widetilde{M}$ consists of the following data:

- A finite set of distinct points $P_1, P_2, \dots, P_n$ in $\widehat{\Sigma}$;
- for each j , a choice of point $Q_j \in F_j = \pi^{-1}(P_j)$;
- for each j , a choice of weight $\alpha_j \in (0, 1) \cap \mathbb{Q}$.

A geometrically ruled surface with a parabolic structure will be called a *parabolic ruled surface*.

If $S \subset \widetilde{M}$ is a holomorphic section of π , we define its slope

$$\mu(S) = S^2 + \sum_{Q_j \notin S} \alpha_j - \sum_{Q_j \in S} \alpha_j;$$

we say that a parabolic ruled surface is *stable* if for every holomorphic section S , we have $\mu(S) > 0$.

3.1. Iterated blow-up of a parabolic ruled surface. Let $\widetilde{M}$ be a parabolic ruled surface. We shall now define a multiple blow-up $\Phi : \widehat{M} \rightarrow \widetilde{M}$ which is canonically determined by the parabolic structure of $\widetilde{M}$.

In order to simplify the notation, suppose that the parabolic structure on $\widetilde{M}$ is reduced to a single point $P \in \widehat{\Sigma}$; let Q be the corresponding point in $F = \pi^{-1}(P)$ and let $\alpha = \frac{p}{q}$ be the weight, where p and q are two coprime integers, $0 < p < q$. Denote the Hirzebruch–Jung continued fraction expansion of α by

$$\frac{p}{q} = \frac{1}{e_1 - \frac{1}{e_2 - \dots - \frac{1}{e_l}}}; \quad (3.1)$$

define also

$$\frac{q-p}{q} = \frac{1}{e'_1 - \frac{1}{e'_2 - \dots - \frac{1}{e'_m}}}. \quad (3.2)$$

These expansions are unique if, as we shall assume, the e_j and e'_j are all ≥ 2 .

We give here a construction the iterated blow-up $\widehat{M}$: the fiber F has self-intersection 0. The first step is to blow up Q , to get a diagram of the form

$$\begin{array}{c} -1 \qquad \qquad -1 \\ \hline \qquad \bigcirc \end{array} \quad (3.3)$$

By blowing up the intersection point of these two curves we get the diagram

$$\begin{array}{ccccccc} & -2 & & -1 & & -2 & \\ & \circ & & \circ & & & \end{array} \quad (3.4)$$

Then we perform an iterated blow-ups of one of the two intersection of the only -1 -curve in the diagram. Given $\alpha = p/q$, there is a unique way (cf. [RS, Proposition 2.1.1]) to choose at each step which point has to be blown-up in order to get the following diagram

$$\begin{array}{cccccccccccccccc} -e_1 & -e_2 & & -e_{l-1} & -e_l & -1 & -e'_m & -e'_{m-1} & & -e'_2 & -e'_1 \\ \circ & \circ & - & \circ & \circ & \circ & \circ & \circ & - & \circ & \circ & & & & \end{array}$$

where the $-e_1$ -curve is the proper transform of the original fiber F .

More generally, if $\widetilde{M}$ has more parabolic points, we perform the same operation for every point and get a corresponding iterated blow-up $\Phi : \widehat{M} \rightarrow \widetilde{M}$.

3.2. The theorem of Mehta-Seshadri. We now give a practical way of using Theorem C. Given a parabolic ruled surface $\widetilde{M} \rightarrow \widehat{\Sigma}$, we deduce a orbifold Riemann surface $\overline{\Sigma}$ by introducing a orbifold singularity of order q_j at every parabolic point $P_j \in \widehat{\Sigma}$ of weight p_j/q_j . As a corollary of Mehta-Seshadri theorem [MS], we have the following proposition.

Proposition 3.2.1. *Let $\widetilde{M} \rightarrow \widehat{\Sigma}$ be a parabolically stable ruled surface. Then there exists an irreducible representation $\rho : \pi_1^{\text{orb}}(\overline{\Sigma}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$ such that $\widehat{M} \simeq \widehat{M}_\rho$, where $\widehat{M}$ is the iterated blow-up of the parabolic ruled surface $\widetilde{M}$ as described in Section 3.1 and $\widehat{M}_\rho$ is the smooth resolution of the orbifold $\overline{M}_\rho$ defined in Section 2.*

Proof. This is a direct consequence of [RS, Theorem 3.3.1]. Q.E.D.

In conclusion we can reformulate Theorem C as follows.

Theorem D. *Let $\widetilde{M} \rightarrow \widehat{\Sigma}$ be a parabolically stable ruled surface, with parabolic weights $\alpha_j = p_j/q_j$ where p_j and q_j are positive coprime integers. Let $\overline{\Sigma}$ be the orbifold ruled surface deduced from $\widehat{\Sigma}$ according to the parabolic structure.*

Then the conclusions (i) and (ii) of Theorem C hold with $\widehat{M}_\rho$ replaced by $\widehat{M}$.

4. THE GOODS: PROOF OF THEOREMS A AND B

4.1. A stable parabolic bundle. Let $\overline{\Sigma}$ be $\mathbb{CP}^1$ with 3 orbifold points P_1, P_2, P_3 , each of weight 2. Then the orbifold Euler characteristic is $1/2$ and $\overline{\Sigma}$ is good, and we can equip $\overline{\Sigma}$ with a metric of CSC $\kappa_1 > 0$. Now, we consider the ruled surface $\pi : \mathbb{CP}^1 \times \mathbb{CP}^1 \rightarrow \mathbb{CP}^1$ where π is, say, the projection on

the first factor. We pick a point Q_j in the fibre $\pi^{-1}(P_j)$, and give it weight $\alpha_j = 1/2$.

Lemma 4.1.1. *For a generic choice of points Q_1, Q_2, Q_3 , the parabolic ruled surface $\widetilde{M} \rightarrow \mathbb{CP}^1$ defined above is parabolically stable.*

Proof. We just need to arrange so that 2 points Q_j never belong to the same constant section of $\widetilde{M} \rightarrow \mathbb{CP}^1$. Q.E.D.

4.2. Conclusion. The iterated blow-up $\widehat{M}$ of Theorem D in this case consists of performing 2 blow-ups in each fibre, so that $\widehat{M}$ is a 6-point blow-up of $\mathbb{CP}^1 \times \mathbb{CP}^1$. Thus this result yields positive-CSC Kähler metrics on X_6 . By adjusting κ_1 and κ_2 and taking the gluing parameter to be small enough, we can get any positive value of c in (1.1).

To deal with X_7 and X_8 , we can use the last part of Theorem C to perform further blow-ups on the previous example. Alternatively, we can replace one of the parabolic weights by $1/3$ or $1/4$ in the above. The complex surface X_8 can also be constructed by adding a fourth parabolic point P_4 , and taking all weights equal to $1/2$. It is readily seen that for generic choices of the points Q_j , the resulting parabolic ruled surfaces are stable.

4.2.1. Remark. — We deliberately chose κ_1 and κ_2 so as to get CSC metrics that are far from being Einstein. However, if we take $\kappa_1 = \kappa_2$ then it seems possible that with a little more work this gluing construction might be adapted to give the KE metrics for (certain complex structures on) X_6 and X_7 ! From this point of view, it would also be interesting to try to extend the analysis to allow $\overline{M}_\rho$ to carry non-trivial holomorphic vector fields. As the analysis in [LSc] shows, it is quite possible for the deformation problem to be unobstructed in this case, although the whole story is then much more delicate.

4.3. Proof of Theorem B. The proof follows the same lines as in [RS] (or as above), taking $\overline{\Sigma}$ to be $\mathbb{CP}^1$ with 4 orbifold points of order 2, 2, 2, 3. The orbifold euler characteristic is $-1/6$ and so we can equip $\overline{M}_\rho$ with a CSC Kähler metric with normalized scalar curvature $c_1 \cdot [\omega] / \sqrt{[\omega]^2}$ equal to any given real number. Applying Theorem D, we get the result claimed in Theorem B.

5. EXPLICIT REPRESENTATIONS

Our use of the theory of stability to generate representations of the orbifold fundamental group results in a certain loss of explicitness—one of the merits of gluing constructions is that they give “nearly” explicit metrics, in the sense that for small values of the gluing parameter, the glued metric is close to the

original one. From this point of view, it seems desirable to try to make explicit these representations, at least in a few examples, and this section is devoted to such a task.

5.1. Platonic orbifold surfaces. Let Γ be one of the finite subgroups of $\mathrm{SO}(3)$ as in Proposition 2.1.2. Then there is a canonical orbifold $\overline{M} = \mathbb{CP}^1 \times \mathbb{CP}^1 / \Gamma$, where Γ acts diagonally on the product. If Γ is not cyclic, then this orbifold will not support any non-trivial holomorphic vector fields and the resolution will carry a positive CSC Kähler metric.

5.2. Representations of the fundamental group of the 4-fold punctured sphere. Let $\overline{\Sigma}$ be the orbifold Riemann surface used in the proof of Theorem B. We shall construct representations of the orbifold fundamental group of $\overline{\Sigma}$. Let the points be denoted P_j , with orders 2, 2, 2, 3 respectively, and let l_j denote the homotopy class of a small loop around P_j . The orbifold fundamental group is

$$\pi_1^{\mathrm{orb}}(\overline{\Sigma}) = \langle l_1, l_2, l_3, l_4 : l_1 l_2 l_3 l_4 = l_1^2 = l_2^2 = l_3^2 = l_4^3 = 1 \rangle.$$

Identifying $\mathrm{SU}(2)$ with the group of unit quaternions, we write down a family of representations $\pi_1^{\mathrm{orb}} \rightarrow \mathrm{SU}(2)/\mathbb{Z}_2$

$$\rho(l_1) = \pm i, \rho(l_2) = \pm j, \rho(l_3) = \pm k \exp(\pi(i \cos \phi + j \sin \phi)/3),$$

$$\rho(l_4) = \pm \exp(-\pi(i \cos \phi + j \sin \phi)/3)$$

Denote by R_1, R_2 and R_3, R_4 the corresponding rotations of $\mathbb{R}^3$. Then R_1, R_2 and R_3 , while R_4 is a $2\pi/3$ -rotation, see Figure 1.

This representation is irreducible and so gives rise to a CSC Kähler orbifold $\overline{M}_\rho$. The corresponding minimal resolution $\widehat{M}_\rho$ is the 9-point blow-up of a minimal ruled surface $\widetilde{M} \rightarrow \mathbb{CP}^1$. We do not know which ruled surface this is for general ϕ , but if $\phi = 0$, we sketch an argument explaining why $\widehat{M}$ is isomorphic to $\mathbb{CP}^1 \times \mathbb{CP}^1$, so that the scalar-flat Kähler manifold $\widehat{M}_\rho$ is isomorphic to a 9-point blow-up of $\mathbb{CP}^1 \times \mathbb{CP}^1$.

Taking $\phi = 0$ from now on, we note that the x -axis is stable under the action of ρ : it is fixed point-wise by R_1 and R_4 and flipped by R_2 and R_3 . Set $Q_1 = (1, 0, 0)$, $Q_2 = (-1, 0, 0)$ and consider the Riemann surface

$$S := (\mathbb{CP}^1 \setminus \{P_j\}) \times_\rho \{Q_1, Q_2\}.$$

This sits in $\overline{M}$, and its closure $\overline{S}$ is a 2-fold ramified cover $p : \overline{S} \rightarrow \overline{\Sigma}$. In $\overline{S}$, there are two orbifold singularities sitting over each of P_1 and P_4 ; on the other hand the cover is ramified over P_2 and P_3 and the corresponding points in $\overline{S}$ are smooth. Thus $\overline{S}$ is an orbifold version of $\mathbb{CP}^1$, with two singular points




FIGURE 1. A representation of the fundamental group of the 4-fold punctured sphere

of weight 2 and two singular points of weight 3. Denote by $\overline{N}$ the associated orbifold ruled surface, and by $\widehat{N}$ its minimal resolution.

Downstairs in $\overline{M}$ there are 8 orbifold singularities, say,

$$A_j, B_j \in \pi^{-1}(P_j).$$

Upstairs in $\overline{N}$, P_1 and P_4 have been double-covered by points $P_1^\pm, P_4^\pm$ and there are accordingly singularities

$$A_1^\pm, B_1^\pm \in \pi^{-1}(P_1^\pm), \quad A_4^\pm, B_4^\pm \in \pi^{-1}(P_4^\pm).$$

The double cover is ramified over P_2 and P_3 , so the points A'_2, A'_3, B'_2, B'_3 corresponding to A_2, A_3, B_2, B_3 are smooth. There is a commutative diagram

$$\begin{array}{ccccc}
 \overline{N} & \xleftarrow{\quad} & \widehat{N} & & \\
 \downarrow p^*\pi & \searrow & \downarrow \psi & & \\
 \overline{S} & \hookrightarrow & \overline{M} & \xleftarrow{\quad} & \widehat{M} \\
 & \searrow p & \downarrow \pi & & \\
 & & \underline{\Sigma} & &
 \end{array}$$

where the dashed arrow is a meromorphic map which becomes a smooth ramified double cover if the points A'_2, A'_3, B'_2, B'_3 are blown up.

The reason for passing to $\overline{N}$ is that it contains two parallel sections s_1 and s_2 , defined by $s_1(Q_1) = Q_1, s_1(Q_2) = Q_2$ and $s_2(Q_1) = Q_2, s_2(Q_2) = Q_1$; let $\overline{S}_j$ denote the closure of $s_j(\overline{S})$ in $\overline{N}$. Because the s_j are parallel, $\overline{S}_j^2 = 0$. By a computation that we omit, the proper transforms $\widehat{S}_j$ of $\overline{S}_j$ in $\widehat{N}$ verify $\widehat{S}_1 \cdot \widehat{S}_2 = 0$ and

$$\widehat{S}_1^2 = \widehat{S}_2^2 = -2,$$

see Figure 2.





FIGURE 2. Configuration of curves in $\widehat{N}$

After a sequence of 5 blow-downs in a suitable order in $\widehat{M}$ (corresponding to 10 blow-downs on $\widehat{N}$) we obtain the configuration of curves shown in Figure 3:

It is a minimal ruled surface with a divisor D verifying $D \cdot F = 1$ and $D^2 = 0$, where F is the class of a generic fibre. Therefore it is isomorphic to $\mathbb{CP}^1 \times \mathbb{CP}^1$. Thus $\widehat{M}$ is a five-point blow-up of the minimal resolution of $(\mathbb{CP}^1 \times \mathbb{CP}^1)/\mathbb{Z}_2$, where the action of $\mathbb{Z}_2$ rotates each factor by an angle π .

FIGURE 3. 10-point blow-down of $\widehat{N}$

It is easy to see that the latter complex surface is a four-point iterated blow-up of $\mathbb{CP}^1 \times \mathbb{CP}^1$. Consider the proper transforms $\widehat{S}_1$ of $\overline{S}_1 = (0 \times \mathbb{CP}^1)/\mathbb{Z}_2$ and $\widehat{S}_2$ of $\overline{S}_2 = (\mathbb{CP}^1 \times 0)/\mathbb{Z}_2$. Then we have $\widehat{S}_1 \cdot \widehat{S}_2 = 0$ and $\widehat{S}_1^2 = \widehat{S}_2^2 = -1$. Hence we have the configuration of curves shown in Figure 4 and after 4 blow-downs in a suitable order, we obtain $\mathbb{CP}^1 \times \mathbb{CP}^1$.

5.3. Representations of the fundamental group of the punctured torus.

Let $\mathbb{T}$ be a compact elliptic curve with one marked point P of weight 2, let $\overline{\Sigma}$ stand for the corresponding orbifold Riemann surface. In terms of standard generators,

$$\pi_1^{\text{orb}}(\overline{\Sigma}) = \langle a, b, l : [a, b]l = l^2 = 1 \rangle.$$

We define a representation $\rho : \pi_1^{\text{orb}}(\overline{\Sigma}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$ as follows

$$\rho(a) = \pm e^{\pi k/4} i, \quad \rho(b) = \pm j, \quad \rho(l) = \pm k.$$

The corresponding rotations R_a, R_b, R_l are shown in in Figure 5.

Denote by $\overline{M}$ the CSC Kähler orbifold ruled surface arising from this representation, by $\widehat{M}$ the minimal resolution, and by $\widetilde{M} \rightarrow \mathbb{T}$ a minimal model for $\widehat{M}$.



FIGURE 4. $\widehat{M}$ with five (-1) -curves blown down

FIGURE 5. A representation of the punctured torus

Then we have

Proposition 5.3.1. *The surface $\widehat{M}$ is a double blow-up of $\widetilde{M}$, and carries a CSC Kähler metric. On the other hand, no smooth blow-down of $\widehat{M}$ admits CSC Kähler metric.*

We give a sketch of the proof of this. The existence of a metric on $\widehat{M}$ is assured by Theorem C. Consider, then, the non-existence statement. The geometrically ruled surface $\widetilde{M}$ cannot admit a CSC Kähler metric for $c_1^2 = 0$,

but $c_1 \neq 0$. In $\widehat{M}$ the only curve that can be blown down smoothly is in the fibre over P . Denote by X this blow-down.

As in the previous example, we can pass to a double cover by remembering the two points on the unit sphere on the axis of R_l . We obtain a double-cover $\overline{N}$ of $\overline{M}$ and a corresponding minimal resolution $\widehat{N}$. By considering the geometry of this double-cover, one can show that the double-cover X' of X is a 2-point blow-up of $\mathbb{P}(L_1 \oplus L_2) \rightarrow \mathbb{T}$, where the points now lie on the sections corresponding to L_1 and L_2 . Such X' does not admit a CSC Kähler metric by [LS, Prop. 3.1], and so X cannot admit a CSC Kähler metric either.

6. HIGHER-DIMENSIONAL EXAMPLES

Since the results of [AP] apply in all dimensions, it is natural to try to extend the foregoing results to higher dimensions. The main problem here is that very little appears to be known about the following basic

Question 6.1. If G is a finite subgroup of U_m acting freely on $\mathbb{C}^m \setminus 0$, and $X \rightarrow \mathbb{C}^m/G$ is a resolution of singularities, does there exist an asymptotically locally euclidean zero CSC Kähler metric on X ?

Joyce [J] proved that if X can be chosen to be a crepant resolution ($c_1(X) = 0$), then X does carry an ALE Ricci-flat metric. In dimensions 2 and 3, this is the case if and only if G (is conjugate to) a subgroup of $SU(m)$. In higher dimensions, however, it is not known in general which singularities $\mathbb{C}^m/G$ admit crepant resolutions.

The other class of examples come from explicit constructions of Kähler metrics on total spaces of line-bundles, as described, for example, in [HS] which contains an extensive survey of the literature. The following result follows easily from the methods of that paper, and is probably well known to many. We sketch the proof in order to make this paper self-contained.

Theorem 6.2. *If $G = \{1, \omega, \dots, \omega^{k-1}\}$, where $\omega = e^{2\pi i/k}$ then there is a resolution X of $\mathbb{C}^{m+1}/G$ that admits an asymptotically locally euclidean zero-CSC Kähler metric.*

Proof. We note that for this group G , we can take X to be the total space of $\mathcal{O}(-k) \rightarrow \mathbb{CP}^m$. Indeed, the total space of $\mathcal{O}(-1)$ can be represented as $U/\mathbb{C}^*$, where

$$U = \{(w, z_0, \dots, z_m) \in \mathbb{C} \times (\mathbb{C}^m \setminus 0)\}$$

the action of $\mathbb{C}^*$ is given by

$$t \cdot (w, z_0, \dots, z_m) = (t^{-1}w, tz_0, \dots, tz_m),$$

and the map

$$\beta(w, z_0, \dots, z_m) \mapsto (wz_0, wz_1, \dots, wz_m)$$

is an invariant description of the blow-up of the origin of $\mathbb{C}^{m+1}$. The action of the group G lifts to the action

$$\omega^r(w, z_0, z_1, \dots, z_m) = (\omega^r w, z_0, z_1, \dots, z_m)$$

on U , with quotient equal to the total space of $\mathcal{O}(-k)$.

Following an Ansatz that goes back to Calabi, we seek a Kähler metric on X of the form

$$\omega_f = \omega_0 + i\partial\bar{\partial}f(t) \quad (6.1)$$

where ω_0 is the standard Kähler form on $\mathbb{CP}^m$ and t is logarithm of the fibre-distance function.

To be more explicit, we work with affine coordinates $(z_1, \dots, z_m)$ on $\mathbb{CP}^m$ so that

$$\omega_0 = (i/2)\partial\bar{\partial}\log(1 + |z_1|^2 + \dots + |z_m|^2).$$

In the same coordinates, the fibre metric on $\mathcal{O}(-k)$ is just given by

$$h^k = (1 + |z_1|^2 + \dots + |z_m|^2)^k.$$

Denote by w a holomorphic fibre coordinate, set $z = \log w$ so that

$$2t = z + \bar{z} + k \log h \quad (6.2)$$

We have

$$i\partial\bar{\partial}t = (ik/2)\partial\bar{\partial}\log h = k\omega_0$$

so that (6.1) can be written

$$\omega_f = (1 + kf'(t))\omega_0 + f''(t)i\partial t \wedge \bar{\partial}t.$$

As explained in [HS], it is better to pass to momentum coordinates, so we introduce

$$\tau = f'(t) \text{ and } \varphi(\tau) = f''(t). \quad (6.3)$$

The advantage of this is that the scalar curvature is given by a very simple formula which we now derive. First of all,

$$\omega_f^{m+1} = (1 + k\tau)^m h^{-m-1} \varphi(\tau) \psi \wedge \bar{\psi} \quad (6.4)$$

where ψ is a holomorphic $(m+1)$ -form, so that

$$\begin{aligned} \rho(\omega_f) &= -i\partial\bar{\partial}\log[(1 + k\tau)^m h^{-m-1} \varphi(\tau)] \\ &= 2(m+1)\omega_0 - i\partial\bar{\partial}\log[(1 + k\tau)^m \varphi(\tau)]. \end{aligned} \quad (6.5)$$

Now for any function of $u(\tau)$,

$$i\partial\bar{\partial}u(\tau) = k\varphi u' \omega_0 + \varphi(\varphi u')' i\partial t \wedge \bar{\partial}t \quad (6.6)$$

where prime denotes differentiation with respect to τ . Applying this with

$$u(\tau) = m \log(1 + k\tau) + \log \varphi(\tau)$$

we obtain

$$\rho(\omega_f) = [2(m+1) - k\varphi u']\omega_0 - \varphi(\varphi u')'i\partial t \wedge \bar{\partial} t. \quad (6.7)$$

The scalar curvature $\sigma(\omega_f)$ is given (up to a factor of 2) by

$$\sigma(\omega_f)\omega_f^{m+1} = (m+1)\rho(\omega_f) \wedge \omega_f^m$$

which leads to the final formula

$$\sigma(\omega_f) = \frac{2m(m+1)}{1+k\tau} - \frac{1}{(1+k\tau)^m} \frac{d^2}{d\tau^2} ((1+k\tau)^m \varphi(\tau)). \quad (6.8)$$

Thus $\sigma(\omega_f) = 0$ if

$$(1+k\tau)^m \varphi(\tau) = \frac{2}{k^2} (1+k\tau)^{m+1} + a\tau + b \quad (6.9)$$

where a and b are constants of integration. The boundary conditions for smoothness of the metric at the zero-section are

$$\varphi(0) = 0, \varphi'(0) = 2 \quad (6.10)$$

which gives

$$\begin{aligned} P(\tau) &:= (1+k\tau)^m \varphi(\tau) \\ &= \frac{2}{k^2} ((1+k\tau)^{m+1} + (k-m-1)(1+k\tau) + m-k). \end{aligned}$$

Geometrically, we have repackaged the complex line-bundle X as the total space of an S^1 -bundle over $\mathbb{CP}^m \times [0, \infty)_\tau$; to check that ω_f is really a metric on this S^1 -bundle we need that $\varphi(\tau) > 0$ for all $\tau \in (0, \infty)$. To see that this is the case, note that

$$P'(\tau) = 0 \text{ if and only if } (1+k\tau)^m = 1 - \frac{k}{m+1}$$

and this latter equation cannot be satisfied if $\tau > 0$. Since $P'(0) = 2$, we see that $P'(\tau) > 0$ for all $\tau > 0$ and since $P(0) = 0$ we have $P(\tau) > 0$ for all $\tau > 0$ as claimed. Thus our zero-CSC Kähler metric on X corresponds to the function

$$\begin{aligned} \varphi(\tau) &= \frac{2}{k^2} \left((1+k\tau) + (k-m-1)(1+k\tau)^{1-m} \right. \\ &\quad \left. + (m-k)(1+k\tau)^{-m} \right). \end{aligned} \quad (6.11)$$

Finally we check the asymptotics of this metric. For this, we return to the blow-up map at the beginning of the proof. In the present coordinates, this is

$$x_0 = w^{1/k}, x_1 = w^{1/k} z_1, \dots, x_m = w^{1/k} z_m, \quad (6.12)$$

where $(x_0, x_1, \dots, x_m)$ are standard linear coordinates on $\mathbb{C}^{m+1}$, and the fractional power of w corresponds to the passage from $\mathcal{O}(-k)$ to its k -fold cover $\mathcal{O}(-1)$. The standard Kähler form on $\mathbb{C}^{m+1}$ is

$$\eta = i\partial\bar{\partial}(|x_0|^2 + \dots + |x_m|^2) = i\partial\bar{\partial}(|w|^{2/k} h) = i\partial\bar{\partial} a e^{2t/k}. \quad (6.13)$$

Thus, this standard Kähler form is of the form (6.1) with

$$f(t) = f_0(t) = e^{2t/k} - t/k. \quad (6.14)$$

This yields

$$1 + k\tau_0 = 2e^{2\tau_0/k}, \quad \varphi_0(\tau_0) = \frac{2}{k^2}(1 + k\tau_0). \quad (6.15)$$

Now since

$$dt = \frac{d\tau_0}{\varphi_0(\tau_0)} = \frac{d\tau}{\varphi(\tau)} \quad (6.16)$$

we see that by suitable choice of integration constants, $\tau - \tau_0$ is order τ_0^{1-m} for large τ_0 , and so

$$f(t) = f_0(t) + O(e^{2(1-m)t/k}) = f_0(t) + O(|x|^{2(1-m)}) \text{ for large } |x|.$$

Hence for large $|x|$,

$$\omega_f = i\partial\bar{\partial}[|x_0|^2 + \dots + |x_m|^2 + O(|x|^{2(1-m)})], \quad (6.17)$$

verifying that ω_f is ALE as required. Q.E.D.

We give two applications of this result:

Theorem 6.3. *Let $S_1, \dots, S_m$ be compact Riemann surfaces, each admitting a holomorphic involution ι_j with isolated fixed points. Suppose further that none of the S_j is $\mathbb{CP}^1$, and that at least one is hyperbolic. Let $\iota = (\iota_1, \dots, \iota_m)$ be the product involution on $S = S_1 \times S_2 \times \dots \times S_m$. Then the minimal resolution $\widehat{M}$ of S/ι admits a negative CSC Kähler metric.*

In addition, any further blow-up of $\widehat{M}$ carries a CSC Kähler metric with negative scalar curvature.

Proof. Equip S_j with a metric of constant curvature κ_j . The product metric on S is not Ricci-flat because there is at least one hyperbolic factor. Moreover the orbifold S/ι has no non-zero holomorphic vector fields. This is clear if each factor is hyperbolic, for then S has no non-zero holomorphic fields. If one or more factor in the product is flat, then M does have non-zero holomorphic vector fields, but none of them is preserved by ι . Hence for any choice $(\kappa_1, \dots, \kappa_m)$, S/ι is an unobstructed CSC Kähler orbifold, and every singularity is modelled on $\mathbb{C}^m/(\pm 1)$. Theorem 6.2, with $k = 2$, now gives a resolution of this singularity, and applying the desingularization theorem of [AP] gives the result. The scalar curvature of $\widehat{M}$ is negative since the sum of the κ_j 's is negative.

For the last statement, the CSC Kähler metric on the blow-up is obtained by gluing the ALE metric provided by Theorem 6.2, with $k = 1$. Q.E.D.

Theorem 6.4. *Let $\overline{\Sigma}$ be a good compact orbifold Riemann surface as explained in Section 2.1, carrying no non-trivial holomorphic vector fields, and with orbifold points of order 2 only. Suppose that $\rho : \pi_1^{\text{orb}}(\overline{\Sigma}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$ is a homomorphism that is irreducible in the sense that the induced action of $\pi_1^{\text{orb}}(\overline{\Sigma})$ fixes no point of $\mathbb{CP}^1$.*

Using the diagonal action of $\pi_1(\overline{\Sigma})$ on $(\mathbb{CP}^1)^m$ induced by ρ , define the twisted product $\overline{M}_\rho = \overline{\Sigma} \times_\rho (\mathbb{CP}^1)^m$. This is a complex orbifold with singularities modelled on $\mathbb{C}^{m+1}/\{\pm 1\}$. Equip $\overline{M}_\rho$ with a twisted product metric of CSC. Then the minimal resolution $\widehat{M}_\rho \rightarrow \overline{M}_\rho$ carries a CSC Kähler metric.

In addition, any further blow-up of $\widehat{M}_\rho$ carries a CSC Kähler metric.

Proof. Again we will use the ALE spaces of Theorem 6.2, with $k = 2$ and $k = 1$ (for the last statement). In order to use the gluing theorem of [AP], the only thing that needs to be checked is that $\overline{M}_\rho$ does not carry any non-trivial holomorphic vector field. But, the argument is a straightforward generalization of the one given [RS] in the case $m = 1$. Q.E.D.

6.4.1. Remark. — In contrast to the previous theorem, by adjusting the curvatures of the base and the fibre here, in the case where $\chi^{\text{orb}}(\overline{\Sigma}) < 0$, we can arrange for the scalar curvature of $\widehat{M}_\rho$ to be positive, zero, or negative.

REFERENCES

- [AP] C. Arezzo, F. Pacard, *Blowing up and desingularizing Kähler orbifolds of constant scalar curvature*, arXiv math.DG/0411522
- [BB] D. Burns, P. de Bartolomeis, *Stability of vector bundles and extremal metrics*, Invent. Math. **92**, 403–407 (1988)

- [CS] D. Calderbank, M. Singer, *Einstein metrics and complex singularities*, Invent. Math. 156 (2004), no. 2, 405–443
- [D] S. Donaldson, *Scalar curvature and projective embeddings I*, J. of Diff. Geom. **59** (2001), 479–522.
- [HS] A.D. Hwang, M.A. Singer, *A Momentum construction for circle-invariant Kähler metrics*, Trans. Amer. Math. Soc. **354** (2002), 2285–2325.
- [J] D.D. Joyce, *Compact manifolds with special holonomy* OUP Oxford (2000).
- [KLP] J. Kim, C. LeBrun, M. Pontecorvo, *Scalar-flat Kähler surfaces of all genera*, J. Reine Angew. Math. **486** (1997), 69–95
- [LS] C. LeBrun, M. Singer, *Existence and deformation theory for scalar-flat Kähler metrics on compact complex surfaces*, Invent. Math. **112** (1993), no. 2, 273–313
- [LSc] C. LeBrun, S.R. Simanca, *Extremal Kähler metrics and complex deformation theory*, Geom. Funct. Anal. 4 (1994), no. 3, 298–336.
- [MS] V.B. Mehta, C.S. Seshadri, *Moduli of vector bundles of curves with parabolic structures*, Math. Ann. **248**, 205–239 (1980)
- [RS] Y. Rollin, M. Singer, *Non minimal scalar-flat Kähler surfaces and parabolic stability*, arXiv math.DG/0404423, to appear in Inventiones Mathematicae.
- [T] G. Tian, *On Calabi’s conjecture for complex surfaces with positive first Chern class* Invent. Math. **101**, no.1, (1990) 101–172.
- [Tr] M. Troyanov, *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc. **324** (1991), no. 2, 793–821

YANN ROLLIN, MIT, 77 MASSACHUSETTS AVENUE, CAMBRIDGE MA 02139, USA
E-mail address: rollin@math.mit.edu

MICHAEL SINGER, SCHOOL OF MATHEMATICS, KING’S BUILDINGS, EDINBURGH SCOTLAND
E-mail address: michael@maths.ed.ac.uk